

2026

Simplified vaccination schedule



Appropriate age	Mandatory vaccinations for infants										6 years	11-13 years	14 years	25 years	65 years +	
	1 month	2 months	3 months	4 months	5 months	6 months	11 months	12 months	16-18 months							
BCG																
DTaP/IPV															Every 20 years	Every 10 years
Hib																
Hepatitis B																
Pneumococcus																
MMR																
Meningococcus ACWY																
Meningococcus B																
Rotavirus																
HPV																
Influenza																Every year
COVID-19																Every year
Shingles																
RSV*																75 years +

*Infants should be immunized against RSV either by vaccination during pregnancy or after birth.

Tuberculosis (BCG)

For children at high risk of tuberculosis, vaccination is recommended from 1 month to 15 years of age.

Diphtheria-Tetanus-Pertussis-Poliomyelitis (DTaP/IPV)

Doses are given at the fixed ages stated in the schedule. Unvaccinated individuals can have catch-up vaccination, given in three doses with a 2-month interval between the first and second dose and a 6-month interval between the second and third dose.

Pregnant women should receive a dose of DTaP/IPV in each pregnancy to protect their infant.

Haemophilus influenzae type b (Hib)

Children who have not been vaccinated can have catch-up vaccination up to the age of 5.

Hepatitis B

If vaccination has not been given during the first year of life, it can be done up to and including the age of 15. From 16 years old, it is recommended only for those exposed to a risk of hepatitis B.

Pneumococcus

Vaccination is mandatory for infants, and catch-up can be given up to 24 months of age. This vaccination is also recommended for all people aged 65 and over.

Measles-Mumps-Rubella (MMR)

People born from 1980 onwards are up to date if they have had two vaccine doses at least one month apart (or three doses if the first dose was received before 12 months).

Meningococcus ACWY

Vaccination is mandatory for infants up to 24 months old. Catch-up can be done up to the age of 5. For young people who were not vaccinated before age 14, catch-up vaccination is recommended between 15 and 24 years old.

Meningococcus B

Vaccination is mandatory for infants up to 24 months old. Catch-up can be done up to the age of 5. For unvaccinated young people, catch-up can be given between the ages of 15 and 24.

Rotavirus

Recommended for all infants from 2 months old. Two or three oral doses are required depending on the vaccine.

Human papillomavirus (HPV)

Both girls and boys aged 11 to 14 should be vaccinated, and catch-up can be given up to and including the age of 26.

Influenza

Vaccination is recommended every year, particularly for individuals with a risk of complications: people aged 65 and over,

people with certain chronic illnesses (including children from 6 months old), pregnant women and people with obesity (BMI > 40 kg/m²). Vaccination against influenza can also be offered to all children aged between 2 and 17 years old.

COVID-19

In the autumn, COVID-19 vaccination is recommended for the same at-risk individuals targeted by the influenza vaccination campaign and people with psychiatric disorders, dementia or Down's syndrome.

In the spring, vaccination is recommended for people aged 80 and over, nursing home residents and patients in long-term care units, and immunocompromised individuals of all ages.

Shingles

Vaccination is recommended for people aged 65 and over. Two doses are given at least two months apart.

RSV

Vaccination is recommended for people aged 75 and over. It is also recommended for people aged 65 and over with chronic respiratory conditions (particularly chronic obstructive pulmonary disease) or heart conditions (particularly heart failure).

Find out more



VACCINATION
INFO SERVICE.FR

Le site de référence qui répond à vos questions

